

The Driving Mechanism Behind Attrition-Enhanced Deracemization**

Wim L. Noorduin, Willem J. P. van Enckevort, Hugo Meekes, Bernard Kaptein, Richard M. Kellogg, John C. Tully, J. Michael McBride, and Elias Vlieg*

The transformation of the enantiomorphic but intrinsically achiral salt NaClO_3 into single chirality was recently demonstrated by Viedma by simply grinding the crystals in a saturated solution.^[1] Broader application of this grinding method to the deracemization of chiral molecules is subject to prerequisites. The enantiomers must crystallize as separate crystals, that is, a racemic conglomerate, and possess the ability to be racemized in solution.^[2] The proof of principle was demonstrated using a derivative of the amino acid phenylglycine.^[3] For both the chiral and nonchiral compounds, the solid phase enriches exponentially.^[4]

Although the practical execution of the process is remarkably simple, there has been much debate regarding its mechanistic details.^[2,4,5-12] There is consensus about the main processes that take place. Crystals of both enantiomorphs are continuously ground to small fragments, part of which dissolve, while at the same time the remaining larger crystals grow until they are crushed again. In a stagnant solution, Ostwald ripening, a near-equilibrium process for which large crystals grow at the cost of smaller ones, leads to one single crystal, and thus of single-handedness.^[4,13] The process slows down extremely when the crystals become larger. During grinding, however, the rate of Ostwald ripening remains high within a steady-state crystal size

distribution that is made up of a large number of small crystals. Several models have demonstrated that for Ostwald ripening to show exponential enantioenrichment in the solid phase, it is not sufficient to describe the crystal growth exclusively by incorporation of monomers;^[4-7] an additional autocatalytic growth mode is necessary to explain the observed exponential behavior. The key question thus lies in the details of the transport of material to the final enantiopure solid phase. Herein this process is discussed for chiral molecules, where solution-phase racemization is necessary.

The first attempt to describe Viedma's experiment in detail was published by Uwaha.^[5a] The autocatalytic behavior in his model is due to the reincorporation of subcritical chiral clusters in crystals of the same handedness. Recently, Saito and Hyuga proposed a physical interpretation in terms of an enhanced racemization reaction at the surface of the enantiomorphic crystals.^[7b] In both models, the minor population is completely converted into the major population of the opposite handedness with a rate that follows the experimentally observed exponential behavior.

The exponential behavior is thus an insufficient criterion to choose one of the models, but, as we will demonstrate below, the models can be distinguished by observing the enantiomeric excess (*ee*) in the solution. Tsogoeva, Mauksch and co-workers have reported enantioenrichment in solution for the enantiomer that forms the minor population in the solid phase during a reversible Mannich reaction.^[11] The authors also conclude that surface-assisted racemization is active in that system.

It was our aim to obtain quantitative data that could be numerically compared with the existing theoretical models. Reliably measuring the solution *ee* during the deracemization is very difficult, and we therefore performed experiments in the absence of racemization for different values of the *ee* in the solid phase. Using a centrifuge, the solids were separated from the liquid phase, thus allowing the measurement of the solution enantiomeric excess as a function of the solid phase *ee* for a wide range of compositions. We found that the solution phase has a surprisingly large excess of molecules with the handedness of the minor solid phase, which becomes more pronounced upon enrichment of the solid phase in the opposite enantiomer. This excess forms the driving force for a net flux of molecules from crystals of the minor handedness to crystals of the major handedness in cases where racemization occurs in solution.

As a model compound we used the same amino acid derivative that led to the proof of principle of the method for chiral molecules, namely the imine formed from 2-methyl-

[*] Dr. W. L. Noorduin, Dr. W. J. P. van Enckevort, Dr. H. Meekes, Prof. Dr. E. Vlieg^[1]
Radboud University Nijmegen
Institute for Molecules and Materials
Heyendaalseweg 135, 6525 AJ Nijmegen (The Netherlands)
Fax: (+31) 24-365-3067
E-mail: e.vlieg@science.ru.nl

Dr. B. Kaptein
DSM Innovative Synthesis BV
PO Box 18, 6160 MD Geleen (The Netherlands)

Prof. Dr. R. M. Kellogg
Syncom BV
Kadijk 3, 9747 AT Groningen (The Netherlands)

Prof. Dr. J. C. Tully, Prof. Dr. J. M. McBride
Department of Chemistry, Yale University
Box 208107, New Haven, CT 06520 (USA)

[†] Present address: WLN: School of Engineering and Applied Sciences, Harvard University
Cambridge, MA (USA)

[**] The SNN agency (Cooperation Northern Netherlands) and the European Fund for Regional Development (EFRO) are acknowledged for partial financial support of this work. The group from Radboud University Nijmegen thanks COST Action CM0703 for support.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201002036>.

benzaldehyde and phenyl glycinamide (**1**).^[3] We started by grinding nonracemic mixtures of **1** (0.26 g) with various values for the initial *ee* in acetonitrile (4.5 g) without racemization catalysts using glass beads (8.5 g) in a thermostated ultrasonic cleaning bath.^[14] After two hours of grinding, 0.8 mL of the slurry was collected and immediately centrifuged at 13 000 rpm for 2 min.^[15] From the upper layer, about 10 μ L of the clear supernatant was carefully removed and analyzed using chiral HPLC. Additionally, the remaining suspension was filtered on a P4 glass filter and the chiral purity of the solid was analyzed using HPLC. Figure 1 shows the solution phase *ee*₁ as a function of the *ee*_s in the solid phase after




Figure 1. Enantiomeric excess in the solution phase as a function of the solid-phase enrichment. The solution enriches in the handedness that is opposite to the enantiomer of the major population in the solid phase. Filled symbols show the results of grinding a solid phase enriched in (*R*)-**1**, leading to a solution phase enriched in (*S*)-**1**. The curve is a fit of Equation (2) to the data ($K_1 = 17.1$, $K_2 = 9.74$).

grinding. As **1** is a racemic conglomerate, thermodynamics predicts that the supernatant solution phase is racemic for any solid composition. We might expect a slight enrichment in the handedness that is abundant in the solid phase if small amounts of solid material are also accidentally sampled. In practice, however, the solution phase is found to be enriched in the enantiomer that forms the minor population in the solid phase. This enantioenrichment becomes even more pronounced the larger the *ee* in the solid phase (Figure 1).

In an alternative experiment to test whether the solution-phase *ee* is indeed the result of the grinding, we waited for four hours between the grinding and centrifuge steps. In this case, the solution-phase enantiomeric excess was zero within the detection limit of the chiral HPLC (ca. 0.5% *ee*). Therefore, the inverted *ee* value in the solution phase is the result of continuous grinding of the enantiomerically enriched conglomerate crystals.

Our results clearly contradict the enhanced surface racemization model of Saito, which predicts a zero solution *ee* without racemization.^[16] To compare the experimental results with the model of Uwaha, we adapted the cluster model to cover the absence of racemization. The crystal size distribution in an actual system is simplified by considering only two sizes: large crystals and small clusters (Figure 2).^[5] The clusters represent the fragments resulting from ablation.^[17] The set of coupled differential equations that describe the development of the total number of molecules in the




Figure 2. Representation of the continuous growth and dissolution of crystals during abrasive grinding, according to McBride.^[6b] During the grinding, crystals are continuously fragmented into clusters. The reincorporation of small clusters occurs more frequently for the major population in the solid phase (process *c*). As a result, a higher fraction of clusters of the minor population dissolve, leading to an increase in molecules of this handedness in the solution phase. Under racemizing conditions, this concentration difference allows a net flux of molecules of the minor population to molecules of the major chirality. Thus, the final result is a complete conversion of the crystals of the minor handedness into crystals of the chirality that initially forms the major population.

(*R*) crystals, B_R , the number in the (*R*) clusters, C_R , and in the solution, M_R , in time t is given by Equations (1a–c):

$$\frac{\partial B_R}{\partial t} = a(M_R - M_{B,eq})B_R - bB_R + cB_R C_R \quad (1a)$$

$$\frac{\partial C_R}{\partial t} = a'(M_R - M_{C,eq})C_R - bB_R - cB_R C_R \quad (1b)$$

$$\frac{\partial M_R}{\partial t} = -a(M_R - M_{B,eq})B_R - a'(M_R - M_{C,eq})C_R \quad (1c)$$

along with analogous equations for the numbers of (*S*) molecules (for a graphical representation, see Figure 2). The kinetic rate constants are a , a' , b , and c , whilst $M_{B,eq}$ and $M_{C,eq}$ are the amounts of solution molecules that would be in equilibrium with crystals and clusters, respectively. The conservation of molecules is covered by Equation (1d):

$$T_R = B_R + C_R + M_R \quad \text{and} \quad T_S = B_S + C_S + M_S \quad (1d)$$

For further details, see Supporting Information. For steady-state conditions, the model yields an analytical expression of the solution phase *ee*₁ in terms of the solid phase *ee*_s [Equation (2)]:

$$ee_1 = \frac{-ee_s}{K_1 - K_2(ee_s)^2} \quad (2)$$

where K_1 and K_2 are fit parameters (see the Supporting Information). A fit of this expression to the experimental data is included in Figure 1; the agreement is excellent.

How can we use these results to understand the grinding-enhanced deracemizations? For that, we need to focus on the cyclic processes that occur during the continuous grinding of

the two populations of crystals (Figure 2). The grinding causes the ablation of small fragments from the crystals. According to the Gibbs–Thomson effect, the small fragments or clusters have a higher solubility than the larger crystals.^[18] As a result, the smallest clusters dissolve, thereby nurturing the larger crystals: Ostwald ripening. The modeling shows that in Equation (1), the parameter c has to be non-zero to achieve an ee in the solution phase as observed in the experiments (see the Supporting Information). This fact implies that there is a direct route from clusters to crystals of the same handedness, irrespective of the actual transfer mechanism of the molecules. Entire clusters could be incorporated, but the results (and mathematics) are the same if, for example, only a fraction of the cluster would transfer to the crystal. These situations correspond to different values of the rate constant c . The only difference between the practical execution of the deracemizations and the results presented herein is the presence or absence of a racemization catalyst. Without racemization, the solution contains an enantiomeric excess that is the inverse of that in the solid phase. In the presence of a racemization catalyst, that is, in a deracemization process, this enantiomeric excess erodes, resulting in an overall net flux of molecules from crystals of the minor handedness towards crystals of the major form. From that perspective, the measurements presented herein can be considered as snapshots of several moments in the deracemization process. In this manner, the inverted enantiomeric excess in the solution phase is the driving force for the deracemization process. Although we have focused on chiral systems, the cluster incorporation model also explains the observed exponential enantiomeric enrichment in achiral systems such as NaClO_3 .^[1]

The solution ee is a consequence of the fact that for the minority solid phase, the clusters have a lower probability of encountering a crystal of the same handedness. This probability can also be lowered by adding more solvent. An interesting and paradoxical consequence is that upon addition of solvent to a ground slurry, the concentration in the solution increases. We indeed observe this phenomenon of reverse dilution (see the Supporting Information). In a deracemization experiment, the initial 50:50 mixture of left- and right-handed solid converts into 100% single chirality. Reverse dilution then leads to a lower concentration in the final solution corresponding to an energetically more favorable state.

Currently, grinding-induced deracemization is being explored as a practical route to produce enantiomerically pure compounds that can be used in the manufacture of pharmaceuticals. It also provides a scenario for the evolution of single chirality as is found in nature.^[12,19–23] Furthermore, a wide variety of procedural approaches, including circularly polarized light, shifted crystal size distributions, and chiral additives have been shown to change at will a racemic mixture of conglomerate crystals to a predetermined handedness.^[3,23–25] Although the rate-determining parameters have been identified, pinning down the driving force for this asymmetric transformation has remained a challenge.^[4,5–12,14] The major question has been why there is a net flux of molecules of the minor handedness in the solid phase towards the major form.

Herein, we have demonstrated experimentally that in the absence of racemization, a concentration inversion in the solution exists during the grinding of a slurry containing an enantioenriched racemic conglomerate. Upon racemization, this solution phase ee drives molecules towards the handedness that already forms the major population in the solid phase.

From these results, we conclude that deracemization of racemic conglomerates can be explained by including four processes: 1) racemization in the solution (not needed for achiral systems), 2) Ostwald ripening; that is, large crystals growing at the expense of smaller crystals, 3) enantioselective (partial) incorporation of clusters, and 4) attrition. The enantioselective cluster incorporation as proposed by Uwaha is a realization of the concept of chiral recognition.^[2a]

There has been some debate as to whether the current process should be called Ostwald ripening or not. It is interesting to note that Liesegang, when he introduced the term Ostwald ripening in 1911, suggested that large crystals could grow from small ones either by coalescence or by transport of material through dissolution, which is equivalent to the route through clusters and monomers, respectively, as shown herein.^[26] Also in a stagnant solution, clusters will be present as a consequence of the dissolution of the smaller crystals, but these will not play an important role during ripening. The continued attrition during grinding experiments, however, results both in a large number of clusters and frequent encounters between clusters and crystals. This factor appears to be essential to explain the exponential increase in solid-phase ee .

The attrition leads to a non-equilibrium crystal size distribution. To stress the difference with the equilibration process of Ostwald ripening, we propose calling this novel route to single chirality Viedma ripening. The reverse dilution forms the thermodynamic driving force in Viedma ripening, whilst the solution-phase racemization provides the kinetic pathway.

Received: April 6, 2010

Revised: June 7, 2010

Published online: September 21, 2010

Keywords: ablation · chiral symmetry breaking · clusters · crystallization · deracemization

-
- [1] C. Viedma, *Phys. Rev. Lett.* **2005**, *94*, 065504.
 [2] a) J. Crusats, S. Veintemillas-Verdaguer, J. M. Ribó, *Chem. Eur. J.* **2006**, *12*, 7776; b) Z. El-Hachemi, J. Crusats, J. M. Ribó, S. Veintemillas-Verdaguer, *Cryst. Growth Des.* **2009**, *9*, 4802.
 [3] W. L. Noorduin, T. Izumi, A. Millemaggi, M. Leeman, H. Meekes, W. J. P. van Enckevort, R. M. Kellogg, B. Kaptein, E. Vlieg, D. G. Blackmond, *J. Am. Chem. Soc.* **2008**, *130*, 1158.
 [4] W. L. Noorduin, H. Meekes, A. A. C. Bode, W. J. P. van Enckevort, B. Kaptein, R. M. Kellogg, E. Vlieg, *Cryst. Growth Des.* **2008**, *8*, 1675.
 [5] a) M. Uwaha, *J. Phys. Soc. Jpn.* **2004**, *73*, 2601; b) M. Uwaha, *J. Phys. Soc. Jpn.* **2008**, *77*, 083802; c) H. Katsuno, M. Uwaha, *J. Cryst. Growth* **2009**, *311*, 4265; d) M. Uwaha, H. Katsuno, *J. Phys. Soc. Jpn.* **2009**, *78*, 023601.

- [6] a) J. M. McBride, J. C. Tully, *Nature* **2008**, *452*, 161; b) J. M. McBride: Lecture at NORDITA Workshop "Origins of homochirality" (February 2008). <http://agenda.albanova.se/conference/Display.py?confId=322>.
- [7] a) Y. Saito, H. Hyuga, *J. Phys. Soc. Jpn.* **2005**, *74*, 535; b) Y. Saito, H. Hyuga, *J. Phys. Soc. Jpn.* **2008**, *77*, 113001.
- [8] a) C. Viedma, *Astrobiology* **2007**, *7*, 312; b) C. Viedma, J. E. Ortiz, T. de Torres, T. Izumi, D. G. Blackmond, *J. Am. Chem. Soc.* **2008**, *130*, 15274.
- [9] D. G. Blackmond, *Chem. Eur. J.* **2007**, *13*, 3290.
- [10] J. H. E. Cartwright, J. M. O. Piro, I. Tuval, *Phys. Rev. Lett.* **2007**, *98*, 165501.
- [11] a) S. B. Tsogoeva, S. Wei, M. Freund, M. Mauksch, *Angew. Chem.* **2009**, *121*, 598; *Angew. Chem. Int. Ed.* **2009**, *48*, 590; b) S. Wei, M. Mauksch, S. B. Tsogoeva, *Chem. Eur. J.* **2009**, *15*, 10255.
- [12] W. L. Noorduin, E. Vlieg, R. M. Kellogg, B. Kaptein, *Angew. Chem.* **2009**, *121*, 9778; *Angew. Chem. Int. Ed.* **2009**, *48*, 9600.
- [13] W. Ostwald, *Z. Phys. Chem.* **1900**, *34*, 495.
- [14] W. L. Noorduin, H. Meekes, W. J. P. van Enkevort, A. Millemaggi, M. Leeman, B. Kaptein, R. M. Kellogg, E. Vlieg, *Angew. Chem.* **2008**, *120*, 6545; *Angew. Chem. Int. Ed.* **2008**, *47*, 6445.
- [15] Similar results were obtained after 16 h of grinding.
- [16] Under racemizing conditions in the solution, surface-assisted racemization might still play a role. The present results indicate that the incorporation of clusters forms the dominant deracemization mechanism, at least in our system.
- [17] In our model, the clusters dissolve and are in that sense subcritical, but they are formed through ablation and not by nucleation. Primary nucleation is unimportant during grinding and is excluded in the model.
- [18] J. W. Gibbs, *Collected Works Vol. I, Thermodynamics*, Yale University Press, New Haven, USA, **1948**.
- [19] W. L. Noorduin, B. Kaptein, H. Meekes, W. J. P. van Enkevort, R. M. Kellogg, E. Vlieg, *Angew. Chem.* **2009**, *121*, 4651; *Angew. Chem. Int. Ed.* **2009**, *48*, 4581.
- [20] M. W. van der Meijden, M. Leeman, E. Gelens, W. L. Noorduin, H. Meekes, W. J. P. van Enkevort, B. Kaptein, E. Vlieg, R. M. Kellogg, *Org. Process Res. Dev.* **2009**, *13*, 1195.
- [21] G. Levilain, C. Rougeot, F. Guillen, J. C. Plaquevent, G. Coquerel, *Tetrahedron: Asymmetry* **2010**, *20*, 2769.
- [22] See the special issue of: *Origins Life Evol. Biospheres* **2010**, *40*, 1.
- [23] W. L. Noorduin, A. A. C. Bode, M. Van der Meijden, H. Meekes, A. F. van Etteer, W. J. P. van Enkevort, P. C. M. Christianen, B. Kaptein, R. M. Kellogg, T. Rasing, E. Vlieg, *Nat. Chem.* **2009**, *1*, 729.
- [24] B. Kaptein, W. L. Noorduin, H. Meekes, W. J. P. van Enkevort, R. M. Kellogg, E. Vlieg, *Angew. Chem.* **2008**, *120*, 7336; *Angew. Chem. Int. Ed.* **2008**, *47*, 7226.
- [25] W. L. Noorduin, W. J. P. van Enkevort, B. Kaptein, R. M. Kellogg, E. Vlieg, *Angew. Chem.* **2010**, *122*, 2593; *Angew. Chem. Int. Ed.* **2010**, *49*, 2539.
- [26] R. E. Liesegang, *Z. Phys. Chem.* **1911**, *34*, 374.